

# Quiver representations: a geometric view\*

Nilava Metya

April 5, 2023

## 1 Representation spaces

Fix a quiver  $Q = (Q_0, Q_1, s, t)$  and a dimension vector  $\mathbf{n} = (n_i)_{i \in Q_0}$ .

**Definition 1** (Representation space). The representation space of the quiver  $Q$  for the dimension vector  $\mathbf{n}$  is

$$\text{Rep}(Q, \mathbf{n}) := \bigoplus_{\{i \rightarrow j\} \in Q_1} \text{Mat}_{n_i \times n_j}(k).$$

This is called the *representation space* because every point  $x \in \text{Rep}(Q, \mathbf{n})$  corresponds to a representation  $V_x$  of  $Q$  with dimension vector  $\mathbf{n}$ . Clearly  $\dim \text{Rep}(Q, \mathbf{n}) = \sum_{\{i \rightarrow j\} \in Q_1} n_i n_j$ . An object  $\mathbf{x} \in \text{Rep}(Q, \mathbf{n})$  will be denoted by  $(x_\alpha)_{(i \rightarrow j) \in Q_1}$  where  $x_\alpha \in \text{Hom}(k^{n_{s(\alpha)}}, k^{n_{t(\alpha)}})$ . The group

$$\text{GL}(\mathbf{n}) := \prod_{i \in Q_0} \text{GL}(n_i)$$

acts on each  $\text{Mat}_{n_i \times n_j}(k)$  by  $(g_i)_{i \in Q_0} \cdot x_\alpha = g_j x_\alpha g_i^{-1}$ , and thus extends to an action on  $\text{Rep}(Q, \mathbf{n})$ . It is not hard to see that  $k^* \cong k^*(\mathbf{1}_{n_i})_{i \in Q_0}$  is a normal subgroup of  $\text{GL}(\mathbf{n})$  and acts trivially on  $\text{Rep}(Q, \mathbf{n})$ . This gives an action of  $\text{PGL}(\mathbf{n}) = \text{GL}(\mathbf{n})/k^*$  on  $\text{Rep}(Q, \mathbf{n})$ . We note that the representations  $V_x, V_y$  for two points  $x, y \in \text{Rep}(Q, \mathbf{n})$  are isomorphic iff  $x, y$  are in the same orbit


---

\*Most material in the notes is basically copied from the mentioned reference article


of  $\mathrm{GL}(\mathbf{n})$  (equivalently,  $\mathrm{PGL}(\mathbf{n})$ ). This is made more formal and informative in the following lemma:

**Lemma 1.1.** *The assignment  $x \mapsto V_x$  gives a one-one correspondence between the orbits  $\mathrm{GL}(\mathbf{n})$  acting on  $\mathrm{Rep}(Q, \mathbf{n})$  and the set of isomorphism classes of representations of  $Q$  with dimension vector  $\mathbf{n}$ . The stabilizer or the isotropy group  $\mathrm{GL}(\mathbf{n})_x = \{g \in \mathrm{GL}(\mathbf{n}) : g \cdot x = x\}$  is isomorphic to the automorphism group  $\mathrm{Aut}_Q(V_x)$ .*

**Example 1.2.** Consider the following quiver



where  $1, n$  denote the dimensions at the respective vertices, so our dimension vector is  $\mathbf{n} = (1, n)$ . Call it  $H_r$ . Then a typical point in  $\mathrm{Rep}(H_r, \mathbf{n})$  looks like  $(M, M_1, \dots, M_r)$  where  $X \in \mathrm{Mat}_{n \times 1}(k) = k^n, M_i \in \mathrm{Mat}_{n \times n}(k)$ . Here  $\mathrm{GL}(\mathbf{n}) = \mathrm{GL}(1) \times \mathrm{GL}(n) = k^* \times \mathrm{GL}(n)$  whose action on  $\mathrm{Rep}(H_r, \mathbf{n})$  is given by  $(c, g) \cdot (M, M_1, \dots, M_r) = (gMt^{-1}, gM_1g^{-1}, \dots, gM_rg^{-1})$ . Such a point corresponds to a representation



The isomorphism classes of representations of the above quiver with the aforementioned dimension vector is parameterized by the orbits of the action of  $\mathrm{GL}(n) = \{1\} \times \mathrm{GL}(n)$  (not just  $\mathrm{GL}(\mathbf{n})$ ) because  $(t, g)$  and  $(1, t^{-1}g)$  have the same action. Basically the action of the  $k^*$  component in  $\mathrm{GL}(\mathbf{n})$  is insignificant in the sense that  $(Mt^{-1}, M_1, \dots, M_n)$  and  $(M, M_1, \dots, M_n)$  belong to the same orbit – we can go from the former to the latter by the action of  $(t^{-1}, \mathrm{Id})$ . Alternately, such a representation is described by a  $k$ -algebra homomorphism  $f : k \langle X_1, \dots, X_n \rangle \rightarrow \mathrm{Mat}_{n \times n}(k), X_i \mapsto M_i$ , together with an element  $M \in k^n$ .

We will call an element  $(M, M_1, \dots, M_r)$  *cyclic* if  $M$  generates  $k^n$  as a  $k \langle X_1, \dots, X_r \rangle$ -module. Collect all such cyclic elements to form the set

$\text{Rep}(H_r, \mathbf{n})^{\text{cyc}}$ . It is clear that  $\text{Rep}(H_r, \mathbf{n})^{\text{cyc}}$  is  $\text{GL}(n)$ -stable. Further if  $\mathbf{M} = (M, M_1, \dots, M_r)$  is a cyclic tuple, then  $\text{GL}(n)_{\mathbf{M}}$  is trivial. This is seen as follows: If  $(\mathbf{M})$  is cyclic, then there are constants  $\lambda_i^{(j)} \in k$  such that  $\sum_i \lambda_i^{(j)} M_i M = \mathbf{e}_j \in k^n$ . If  $g \in \text{GL}(n)_{\mathbf{M}}$  then  $g\mathbf{e}_j = \sum_i \lambda_i^{(j)} gM_i M = \sum_i \lambda_i^{(j)} M_i gM = \sum_i \lambda_i^{(j)} M_i M = \mathbf{e}_j$ . Since this is true for every coordinate vector, we must have  $g = \text{Id}$ .

Let's talk about the orbit space  $\text{Rep}(H_r, \mathbf{n})^{\text{cyc}} / \text{GL}(n)$ . Here we will view the points of representation space as an algebra homomorphism  $k \langle X_1, \dots, X_r \rangle \rightarrow \text{Mat}_{n \times n}(k)$  together with an element of  $k^n$ . This means  $\text{Rep}(H_r, \mathbf{n})^{\text{cyc}} = \{(f, v) \in \text{Hom}(k \langle X_1, \dots, X_r \rangle, \text{Mat}_{n \times n}(k)) \times k^n : f(k \langle X_i \rangle) v = k^n\}$ . Note that Two points  $(M, \mu), (N, \nu) \in \text{Rep}(H_r, \mathbf{n})^{\text{cyc}}$  are in the same orbit iff  $M = gN$  and  $\mu(X_i) = g\nu(X_i)g^{-1}$  for some  $g \in \text{GL}(n)$ . Just to repeat,  $\mu, \nu$  are algebra homomorphisms of the above type. Given any  $(f, v) \in \text{Rep}(H_r, \mathbf{n})^{\text{cyc}}$ , we can put a ring structure on  $k^n$  given as follows: for  $u_1, u_2 \in k^n$  there are polynomials  $P_1, P_2 \in k \langle X_1, \dots, X_r \rangle$  such that  $f(P_i)v = u_i$ , and so define  $u_1 u_2 = f(P_1 P_2)v$ .<sup>1</sup> The kernel is  $I(f, v) = \{P \in k \langle X_1, \dots, X_n \rangle : f(P)v = 0\}$ . One should check that the following is a bijective correspondence between  $\text{Rep}(H_r, \mathbf{n})^{\text{cyc}} / \text{GL}(n)$  and  $\{\text{left ideals} \subseteq k[X_1, \dots, X_n] \text{ of codimension } n\}$ :

$$(f, v) \mapsto I(f, v)$$

$$(P \mapsto (\pi(Q) \mapsto \pi(PQ)), \pi(1)) \leftarrow I$$

$$PQ - P'Q = PQ - PQ' + PQ' - P'Q = PQ' - P'Q$$

**Definition 2.** An *(affine) algebraic group* is an (affine) algebraic variety  $G$  equipped with a group structure such that the multiplication map  $G \times G \rightarrow G$  and the inverse map  $G \rightarrow G$  are morphisms of varieties.

An *algebraic action* of an algebraic group  $G$  on a variety  $X$  is a group action  $G \times X \rightarrow X$  which is also a morphism of varieties.

**Proposition 1.3.** *Let  $G$  have an algebraic action on a variety  $X$ . Fix  $x \in X$ .*

- (a)  $G_x = \{g \in G : g \cdot x = x\}$  is closed in  $G$ .
- (b)  $G \cdot x$  is a locally closed, non-singular subvariety of  $X$ . All connected components of  $G \cdot x$  have dimension  $\dim G - \dim G_x$ .

---

<sup>1</sup>I couldn't verify that this is well defined because of the non-commutativity of the  $X_i$ 's.

- (c) The orbit closure  $\overline{G \cdot x}$  is the union of  $G \cdot x$  and of orbits of smaller dimension; it contains at least one closed orbit.
- (d) The variety  $G$  is connected if and only if it is irreducible; then the orbit  $G \cdot x$  and its closure are irreducible as well.

Now consider a group homomorphism  $\varphi : G \rightarrow H$  of algebraic groups. This gives an action of  $G$  on  $H$  given by  $g \cdot h := \varphi(g)h$ . This is an algebraic action and its orbits are  $G \cdot h = (\text{Im } \varphi) \cdot h$ . There is at least one closed orbit (contained in  $\overline{(\text{Im } \varphi) \cdot h}$  for some  $h \in H$ ). But the orbits are permuted transitively by the action of  $H$  on itself by right multiplication, thus implying that all orbits (that is, cosets) are closed. This means  $\text{Im } \varphi$  is closed. Now note that  $G_{1_H} = \{g \in G : \varphi(g) = 1\} = \ker \varphi$ , which is also closed. Thus  $\ker \varphi, \text{Im } \varphi$  are closed in  $G, H$  respectively. Finally we get that  $\dim \text{Im } \varphi = \dim(G \cdot 1_H) = \dim G - \dim G_{1_H} = \dim G - \dim \ker \varphi$ .

## 2 Isotropy groups

**Proposition 2.1.** *Let  $M$  be a finite-dimensional representation of  $Q$ .*

- (a) *The automorphism group  $\text{Aut}_Q(M)$  is an open affine subset of  $\text{End}_Q(M)$ . As a consequence,  $\text{Aut}_Q(M)$  is a connected linear algebraic group.*
- (b) *There exists a decomposition  $\text{Aut}_Q(M) \cong U \times \prod_{i=1}^r \text{GL}(m_i)$  where  $U$  is a closed normal unipotent subgroup and  $m_1, \dots, m_r$  denote the multiplicities of the indecomposable summands of  $M$ .*

We will allude to a theorem for finite-dimensional representations of associative algebras, and leave it as an exercise to the reader to prove proposition 2.1.

**Theorem 2.2.** *Let  $M$  be a finite-dimensional module over an algebra  $A$ . Then there is a decomposition of  $A$ -modules*

$$M \cong \bigoplus_{i=1}^r M_i^{m_i}$$

where  $M_1, \dots, M_r$  are indecomposable and pairwise non-isomorphic, and  $m_1, \dots, m_r$  are positive integers. Moreover, the indecomposable summands

$M_i$  and their multiplicities  $m_i$  are uniquely determined up to reordering. We also have a decomposition of vector spaces

$$\text{End}_A(M) \cong I \oplus \prod_{i=1}^r \text{Mat}_{m_i \times m_i}(k)$$

where  $I$  is a nilpotent ideal.

*Proof sketch of proposition 2.1.* The first part is immediate by the observation that  $\text{Aut}_Q(M) = \text{End}_Q(M) \setminus V(\det) = D(\det)$ .

For the next part, we start with the split surjective algebra-homomorphism  $\text{End}_Q(M) \rightarrow \text{End}_Q(M)/I \cong \prod_{i=1}^r \text{Mat}_{m_i \times m_i}(k)$  which, in turn, gives a split

surjective algebra-homomorphism  $\text{Aut}_Q(M) \rightarrow \prod_{i=1}^r \text{GL}(m_i)$ . The kernel of this map is  $\text{Id}_M + I$ . Thus,  $\text{Id}_M + I$  is a closed connected normal subgroup of  $\text{Aut}_Q(M)$ .

Next consider the linear action of  $\text{Id}_M + I$  on  $k \text{Id}_M \oplus I$  by left multiplication. Since the orbit of  $\text{Id}_M$  is isomorphic to the affine space  $\text{Id}_M + I$ , this action yields a closed embedding  $\text{Id}_M + I \hookrightarrow \text{GL}(k \text{Id}_M \oplus I)$ . The powers  $I^n$  form a decreasing filtration of the vector space  $k \text{Id}_M \oplus I$ , and they stabilize to 0. Any  $I^n$  is stable under the action of  $I + \text{Id}_M$  and this action fixes the associated grades  $I^n/I^{n+1}$  and the quotient  $(k \text{Id}_M \oplus I)/I$ . This establishes  $\text{Id}_M + I$  as a unipotent subgroup of  $\text{GL}(k \text{Id}_M \oplus I)$ , by choosing a basis of  $k \text{Id}_M \oplus I$  compatible with the filtration  $(I^n)_{n \geq 1}$ . ■

**Corollary 2.3.** *The representation  $V_x$ , for  $x \in \text{Rep}(Q, \mathbf{n})$  is indecomposable if and only if the isotropy group  $\text{GL}(\mathbf{n})_x$  is the semi-direct product of a unipotent subgroup with the group  $k^* \text{Id}_{\mathbf{n}}$ ; equivalently,  $\text{PGL}(\mathbf{n})_x$  is unipotent.*

Now, when studying homological aspects, one comes across the following exact sequence

$$0 \rightarrow \text{End}_Q(M) \rightarrow \prod_{i \in Q_0} \text{End}(V_i) \rightarrow \prod_{\alpha \in Q_1} \text{Hom}(V_{s(\alpha)}, V_{t(\alpha)}) \rightarrow \text{Ext}_Q^1(M, M) \rightarrow 0.$$

The above discussion helps put this exact sequence in a nice geometric framework.

**Theorem 2.4.** *Let  $x \in \text{Rep}(Q, \mathbf{n})$  and denote by  $M = V_x$  the corresponding representation of  $Q$ .*

(a) *There is an exact sequence*

$$0 \longrightarrow \text{End}_Q(M) \longrightarrow \text{End}(\mathbf{n}) \xrightarrow{c_x} \text{Rep}(Q, \mathbf{n}) \longrightarrow \text{Ext}_Q^1(M, M) \longrightarrow 0$$

$$\text{with } c_x((f_i)_{i \in Q_0}) = (f_{t(\alpha)}x_\alpha - x_\alpha f_{s(\alpha)})_\alpha.$$

(b)  *$c_x$  may be identified with the differential at the identity of the orbit map  $\varphi_x : \text{GL}(\mathbf{n}) \rightarrow \text{Rep}(Q, \mathbf{n}), g \mapsto g \cdot x$ .*

(c) *The image of  $c_x$  is the Zariski tangent space  $T_x(\text{GL}(\mathbf{n}) \cdot x)$  viewed as a subspace of  $T_x(\text{Rep}(Q, \mathbf{n})) \cong \text{Rep}(Q, \mathbf{n})$ .*

*Proof.*(b)  $\text{GL}(\mathbf{n}) \subset_{\text{open}} \text{End}(\mathbf{n})$ . So the Zariski tangent space<sup>2</sup> to this group at  $\text{Id}_{\mathbf{n}}$  may be identified with the vector space  $\text{End}(\mathbf{n})$ . The tangent space to  $\text{Aut}_Q(M)$  at  $\text{Id}_{\mathbf{n}}$  is  $\text{End}_Q(M)$ . The action of  $\text{GL}(\mathbf{n})$  is given by

$$\begin{aligned} \text{GL}(n_i) \times \text{GL}(n_j) &\longrightarrow \text{Mat}_{n_i \times n_j}(k) \\ (g, h) &\longmapsto hx_{i \rightarrow j}g^{-1} \end{aligned}$$

$c_x$  immediately comes from the differential of this map

$$\begin{aligned} \text{Mat}_{n_i \times n_i} \times \text{Mat}_{n_j \times n_j} &\longrightarrow \text{Mat}_{n_i \times n_j}(k) \\ (f_i, f_j) &\longmapsto f_j x_{i \rightarrow j} - x_{i \rightarrow j} f_i. \end{aligned}$$

(c) From proposition 1.3 we get  $\dim(\text{GL}(\mathbf{n}) \cdot x) = \dim \text{GL}(\mathbf{n}) - \dim \text{GL}(\mathbf{n})_x$ . But  $\dim(\text{GL}(\mathbf{n}) \cdot x) = \dim [T_x(\text{GL}(\mathbf{n}) \cdot x)]$ . But  $\text{GL}(\mathbf{n})_x$  comprise of the invertible intertwiners for the module  $M = V_x$ , and thus  $\dim \text{GL}(\mathbf{n})_x = \dim \text{Aut}_Q(M) = \dim \text{End}_Q(M)$ . Also  $\dim \text{GL}(\mathbf{n}) = \dim \text{End}(\mathbf{n})$ . The last two equalities follow from the fact that  $\text{GL} \subset_{\text{open}} \text{End}$  which is proposition 2.1. Combining these gives  $\dim [T_x(\text{GL}(\mathbf{n}) \cdot x)] = \dim \text{End}(\mathbf{n}) - \dim \text{End}_Q(M)$ . By (b) and the above exact sequence,  $\ker c_x = \text{End}_Q(M)$ . This means that  $T_x(\text{GL}(\mathbf{n}) \cdot x)$  is the entire image of  $c_x$ . ■

---

<sup>2</sup>This is a technical term which can be defined without using differential geometry concepts and simply by linearizing things using ‘abstract’ algebra. A similar definition in this spirit is the tangent space for a local ring  $(R, \mathfrak{m})$  which is  $\mathfrak{m}/\mathfrak{m}^2$  – this essentially keeps only linear terms.

## References

- [1] M. Brion, “Representations of quivers,” 2008. [Online]. Available: [https://www-fourier.ujf-grenoble.fr/~mbrion/notes\\_quivers\\_rev.pdf](https://www-fourier.ujf-grenoble.fr/~mbrion/notes_quivers_rev.pdf)